



NOAA
FISHERIES

SEFSC, Miami

SEDAR 61: US Gulf of Mexico Red Grouper Interim Analysis

Gulf of Mexico Fishery Management Council
Scientific and Statistical Committee




January 9, 2020

**NOTE: This presentation modified to
focus only on the Interim Analysis**

Interim Analysis

Interim Assessment (IA)

- First red grouper IA conducted in October 2018
- Updated red grouper IA to adjust harvest recommendations based on current stock conditions
 - SEDAR61 terminal year: 2017
 - Made assumptions in projections regarding the impact of 2018 red tide







Cause for Concern

Commercial data from 2010 through 2018 were obtained from the IFQ Quotas and Catch Allowances, accessed December 4, 2019

(<https://portal.southeast.fisheries.noaa.gov/reports/cs/CommercialQuotasCatchAllowanceTable.pdf>), remaining years were obtained from the Gulf of Mexico Historical Commercial Landings and Annual Catch Limits (ACLs), updated November 7, 2018 (<https://www.fisheries.noaa.gov/southeast/gulf-mexico-historical-commercial-landings-and-annual-catch-limit-monitoring/gulf-commercial-historical.pdf>).


Recreational data from 2010 through 2017 were obtained from recreational historical landings, updated March 9, 2019 (<https://www.fisheries.noaa.gov/southeast/recreational-fishing-data/gulf-mexico-historical-recreational-landings-and-annual-catch>), data from 2018 and 2019 (through June) were obtained December 4, 2019 from

<https://www.fisheries.noaa.gov/southeast/2018-and-2019-gulf-mexico-recreational-landings-and-annual-catch-limits-acls-and-annual>.



Index of abundance: NMFS Bottom Longline

- Compare where we are now to where we want to be
 - Where we are now = Observed index value
 - Where we want to be = Forecasted index value
 - Good agreement between values



Harvest Control Rule (HCR)

$$ABC_y = ABC_{assess} \left(\frac{O_y + \beta}{F_y + \beta} \right)$$

- Where $ABC_{assess} = 4.9^*$ million pounds gutted weight

*Sep 2019 SSC recommendation
(76%com,24%rec)

O_y = Observed index value in year y,

F_y = Forecasted index value in year y,

β = Scalar to adjust responsiveness of HCR x SE
of index


- Changes to ABC only implemented every other year

$ABC_{assess} = 4.9^*$ million pounds gutted weight
 *Sep 2019 SSC recommendation (76%com,24%rec)

Effect of β

- Previously recommended: (1) excluding ABC change following SEDAR42; and (2) $\beta = 1$
 - Low β tracks index more closely
 - High β tracks ABC

β	Adj ABC (million pounds gutted weight)
1	5.097
3	5.004
5	4.971
7	4.954
9	4.943



The dashed black line indicates ignoring the ABC increase that resulted from SEDAR42

Ongoing Work:

- Finish MSE development
 - MSE will be used to select HCR/index pair that best achieves management goals
- Design multitude of HCR's with stakeholder input
- Test HCR/Index combinations to identify optimum HCR

Questions?

**Thank you to all SEDAR61 data providers
and for your attention!**

NMFS Bottom Longline Index

Survey Year	Frequency	N	Delta-Lognormal Index	Scaled Index (mean = 1)	Coefficient of Variation	Lower Confidence Level	Upper Confidence Level
2001	0.215	93	0.764	0.827	0.290	0.468	1.460
2002							
2003	0.342	117	1.005	1.088	0.202	0.729	1.624
2004	0.418	98	1.625	1.760	0.193	1.201	2.578
2005	0.250	40	0.559	0.606	0.407	0.277	1.326
2006	0.282	39	0.530	0.574	0.392	0.269	1.224
2007	0.190	42	0.842	0.911	0.465	0.376	2.208
2008	0.267	60	0.582	0.630	0.323	0.336	1.182
2009	0.349	63	0.897	0.971	0.264	0.578	1.633
2010	0.328	67	1.231	1.333	0.266	0.790	2.248
2011	0.402	122	2.310	2.501	0.181	1.745	3.585
2012	0.469	49	2.110	2.285	0.255	1.384	3.772
2013	0.340	47	0.979	1.060	0.306	0.583	1.927
2014	0.262	42	0.568	0.615	0.383	0.293	1.289
2015	0.245	53	0.706	0.764	0.361	0.379	1.540
2016	0.184	49	0.339	0.367	0.436	0.159	0.846
2017	0.318	44	0.701	0.759	0.342	0.390	1.477
2018	0.184	49	0.428	0.463	0.427	0.204	1.050
2019	0.205	39	0.448	0.486	0.461	0.202	1.167