



*With Great Power Comes Great Responsibility: 10 Things
to Know About Steepness for Stock Assessments*

Marc Mangel

Department of Applied Mathematics, UCSC

Department of Biology, University of Bergen

Puget Sound Institute, University of Washington, Tacoma

JACK BASKIN SCHOOL OF ENGINEERING

BIOTECHNOLOGY, INFORMATION TECHNOLOGY, NANOTECHNOLOGY

Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability



Ray (left) and Sidney, 1949

Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$



Ray (left) and Sidney, 1949

Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability



Ray (left) and Sidney, 1949

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$

Comment: What follows can be done for Ricker SRR if that is your preference

Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability



Ray (left) and Sidney, 1949

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$

α = Recruits as eggs become infinite


Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability



Ray (left) and Sidney, 1949

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$

α = Recruits as eggs become infinite



Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability



Ray (left) and Sidney, 1949

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$

α = Recruits as eggs become infinite

$\frac{\alpha}{\beta}$ = Maximum egg survival

Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$

α = Recruits as eggs become infinite

$\frac{\alpha}{\beta}$ = Maximum egg survival



Ray (left) and Sidney, 1949



Jon, Dick Gomulkiewicz, Marc 1989

Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability



Ray (left) and Sidney, 1949

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$

α = Recruits as eggs become infinite

$\frac{\alpha}{\beta}$ = Maximum egg survival



Jon, Dick Gomulkiewicz, Marc 1989

$$R(E_T) = \frac{\alpha E_T}{1 + \beta E_T}$$

Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability



Ray (left) and Sidney, 1949

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$

α = Recruits as eggs become infinite

$\frac{\alpha}{\beta}$ = Maximum egg survival



Jon, Dick Gomulkiewicz, Marc 1989

$$R(E_T) = \frac{\alpha E_T}{1 + \beta E_T}$$

α = Maximum egg survival

Thing #1: How We Write the Beverton-Holt SRR Affects Interpretability



Ray (left) and Sidney, 1949

$$R(E_T) = \frac{\alpha E_T}{\beta + E_T}$$

α = Recruits as eggs become infinite

$\frac{\alpha}{\beta}$ = Maximum egg survival



Jon, Marc 2010

$$R(E_T) = \frac{\alpha E_T}{1 + \beta E_T}$$

α = Maximum egg survival

$\frac{1}{\beta}$ = Egg numbers giving half of
maximum egg survival

Thing #2: Maximum Per Egg Survival Depends on Early Life History

Thing #2: Maximum Per Egg Survival Depends on Early Life History

α = Maximum egg survival

Thing #2: Maximum Per Egg Survival Depends on Early Life History

α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

Thing #2: Maximum Per Egg Survival Depends on Early Life History

α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

Thing #2: Maximum Per Egg Survival Depends on Early Life History

α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

$$\alpha = s_{ER}(\tau_{ER})$$

Thing #2: Maximum Per Egg Survival Depends on Early Life History

α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

$$\alpha = s_{ER}(\tau_{ER})$$

McGurk (1986 MEPS 34:227) Shows How to Find Egg to Recruit Survival

Thing #2: Maximum Per Egg Survival Depends on Early Life History


α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

$$\alpha = s_{ER}(\tau_{ER})$$

McGurk (1986 MEPS 34:227) Shows How to Find Egg to Recruit Survival



Thing #2: Maximum Per Egg Survival Depends on Early Life History


α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

$$\alpha = s_{ER}(\tau_{ER})$$

McGurk (1986 MEPS 34:227) Shows How to Find Egg to Recruit Survival



$$\frac{d}{dt}s_{ER} = -m(w_{ER})s_{ER}$$

Thing #2: Maximum Per Egg Survival Depends on Early Life History


α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

$$\alpha = s_{ER}(\tau_{ER})$$

McGurk (1986 MEPS 34:227) Shows How to Find Egg to Recruit Survival



$$\frac{d}{dt}s_{ER} = -m(w_{ER})s_{ER}$$

If ER Interval Growth is Exponential

Thing #2: Maximum Per Egg Survival Depends on Early Life History


α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

$$\alpha = s_{ER}(\tau_{ER})$$

McGurk (1986 MEPS 34:227) Shows How to Find Egg to Recruit Survival



$$\frac{d}{dt}s_{ER} = -m(w_{ER})s_{ER}$$

If ER Interval Growth is Exponential

$$w_{ER}(\tau) = w_{ER}(0)e^{c_{ER}\tau}$$

Thing #2: Maximum Per Egg Survival Depends on Early Life History


α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

$$\alpha = s_{ER}(\tau_{ER})$$

McGurk (1986 MEPS 34:227) Shows How to Find Egg to Recruit Survival



$$\frac{d}{dt}s_{ER} = -m(w_{ER})s_{ER}$$

If ER Interval Growth is Exponential

$$w_{ER}(\tau) = w_{ER}(0)e^{c_{ER}\tau}$$

Thing #2: Maximum Per Egg Survival Depends on Early Life History


α = Maximum egg survival

$s_{ER}(\tau)$ = Egg survival to time τ of the ER interval

τ_{ER} = Length of the egg to recruit interval

$$\alpha = s_{ER}(\tau_{ER})$$

McGurk (1986 MEPS 34:227) Shows How to Find Egg to Recruit Survival



$$\frac{d}{dt}s_{ER} = -m(w_{ER})s_{ER}$$

If ER Interval Growth is Exponential


$$w_{ER}(\tau) = w_{ER}(0)e^{c_{ER}\tau}$$

- Initial egg/larval size
- Size at recruitment

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data



*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data { $A_{max} = \text{Maximum age}$

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data { $A_{max} = \text{Maximum age}$
 $S(a) = \text{Pr}\{\text{Survive to age } a\}$

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data {
 A_{max} = Maximum age
 $S(a)$ = Pr{Survive to age a }
 $L(a)$ = Length at age a

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data {

- A_{max} = Maximum age
- $S(a)$ = $\Pr\{\text{Survive to age } a\}$
- $L(a)$ = Length at age a
- $p_m(a)$ = $\Pr\{\text{mature at age } a\}$

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data {

- A_{max} = Maximum age
- $S(a)$ = Pr{Survive to age a }
- $L(a)$ = Length at age a
- $p_m(a)$ = Pr{mature at age a }
- b = Length fecundity exponent

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data {

- A_{max} = Maximum age
- $S(a)$ = Pr{Survive to age a }
- $L(a)$ = Length at age a
- $p_m(a)$ = Pr{mature at age a }
- b = Length fecundity exponent
- $\phi(a)$ = Specific fecundity at age a

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data {

- A_{max} = Maximum age
- $S(a)$ = Pr{Survive to age a }
- $L(a)$ = Length at age a
- $p_m(a)$ = Pr{mature at age a }
- b = Length fecundity exponent
- $\phi(a)$ = Specific fecundity at age a

Here's steepness (Mangel et al 2010, 2013)

*Thing #3: As Soon As You Have a Demographic Data for the Stock You
Have a Point Estimate for Steepness*

Demographic data {

- A_{max} = Maximum age
- $S(a)$ = Pr{Survive to age a }
- $L(a)$ = Length at age a
- $p_m(a)$ = Pr{mature at age a }
- b = Length fecundity exponent
- $\phi(a)$ = Specific fecundity at age a

Here's steepness (Mangel et al 2010, 2013)

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}$$

Thing #3: As Soon As You Have a Demographic Data for the Stock You Have a Point Estimate for Steepness

Demographic data {

- A_{max} = Maximum age
- $S(a)$ = Pr{Survive to age a }
- $L(a)$ = Length at age a
- $p_m(a)$ = Pr{mature at age a }
- b = Length fecundity exponent
- $\phi(a)$ = Specific fecundity at age a

Here's steepness (Mangel et al 2010, 2013)

Life time individual egg production

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}$$


*Thing #4: We Can Use Demographic Information and Simulation to
Obtain a Prior for Steepness*

Thing #4: We Can Use Demographic Information and Simulation to Obtain a Prior for Steepness

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}$$

Thing #4: We Can Use Demographic Information and Simulation to Obtain a Prior for Steepness


$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}$$




Bluefin tuna
Mangel et al FaF (2010)

Thing #4: We Can Use Demographic Information and Simulation to Obtain a Prior for Steepness

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a)L(a)^b p_m(a)\phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a)L(a)^b p_m(a)\phi(a)}$$




Bluefin tuna
Mangel et al FaF (2010)




Striped Marlin
Brodziak et al Fish Res 2014

Thing #4: We Can Use Demographic Information and Simulation to Obtain a Prior for Steepness


$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a)L(a)^b p_m(a)\phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a)L(a)^b p_m(a)\phi(a)}$$



Bluefin tuna
Mangel et al FaF (2010)



Striped Marlin
Brodziak et al Fish Res 2014



Blue shark
Kai and Fujinami Fish. Res. 2017

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}$$

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}$$


Write out survival

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}$$

Write out survival

Rate of mortality at age




$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} \prod_{a'=0}^{a-1} e^{-M(a')} L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} \prod_{a'=0}^{a-1} e^{-M(a')} L(a)^b p_m(a) \phi(a)}$$

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} S(a) L(a)^b p_m(a) \phi(a)}$$

Write out survival

Rate of mortality at age



$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} \prod_{a'=0}^{a-1} e^{-M(a')} L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} \prod_{a'=0}^{a-1} e^{-M(a')} L(a)^b p_m(a) \phi(a)}$$

Case of constant mortality: $M(a) = M$

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse


Case of Constant Mortality

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}$$

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

Case of Constant Mortality

Okay to fix this




$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}$$

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

Case of Constant Mortality


Or this


$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}$$

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

Case of Constant Mortality

But both?


$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}$$

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

Case of Constant Mortality

But both?

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}$$



Best result

Thing #5: Specifying Both Natural Mortality and Steepness Will Lead to Problems, Or Worse

Case of Constant Mortality

But both?

$$h = \frac{\alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}{4 + \alpha \cdot \sum_{a=0}^{A_{max}} e^{-Ma} L(a)^b p_m(a) \phi(a)}$$



Best result



Worst result

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - \textcircled{M} N$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN \quad \text{set} \quad E_T = \phi N \quad \text{and} \quad \begin{aligned} \alpha_p &= \alpha \phi \\ \beta_p &= \beta \phi \end{aligned}$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN \quad \text{set } E_T = \phi N \quad \text{and} \quad \begin{aligned} \alpha_p &= \alpha \phi \\ \beta_p &= \beta \phi \end{aligned}$$

$$\frac{dN}{dt} = \frac{\alpha_p N}{1 + \beta_p N} - MN$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN \quad \text{set } E_T = \phi N \quad \text{and} \quad \begin{aligned} \alpha_p &= \alpha \phi \\ \beta_p &= \beta \phi \end{aligned}$$

$$\frac{dN}{dt} = \frac{\alpha_p N}{1 + \beta_p N} - MN$$

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right)$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN \quad \text{set } E_T = \phi N \quad \text{and} \quad \begin{aligned} \alpha_p &= \alpha \phi \\ \beta_p &= \beta \phi \end{aligned}$$

$$\frac{dN}{dt} = \frac{\alpha_p N}{1 + \beta_p N} - MN$$

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right)$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN \quad \text{set } E_T = \phi N \quad \text{and} \quad \begin{aligned} \alpha_p &= \alpha \phi \\ \beta_p &= \beta \phi \end{aligned}$$

$$\frac{dN}{dt} = \frac{\alpha_p N}{1 + \beta_p N} - MN$$

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right)$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN \quad \text{set } E_T = \phi N \quad \text{and} \quad \begin{aligned} \alpha_p &= \alpha \phi \\ \beta_p &= \beta \phi \end{aligned}$$

$$\frac{dN}{dt} = \frac{\alpha_p N}{1 + \beta_p N} - MN$$

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right) \quad \begin{array}{l} \text{Reason for writing the} \\ \text{BH-SRR as above} \end{array}$$

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN \quad \text{set } E_T = \phi N \quad \text{and} \quad \begin{aligned} \alpha_p &= \alpha \phi \\ \beta_p &= \beta \phi \end{aligned}$$

$$\frac{dN}{dt} = \frac{\alpha_p N}{1 + \beta_p N} - MN$$

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right) \quad \begin{array}{l} \text{Reason for writing the} \\ \text{BH-SRR as above} \end{array}$$

Here's steepness (Mangel et al 2010, 2013)

Thing #6: Although They Are Not Fashionable Just Now, Production

Models Can Teach Us a LOT

$$\frac{dN}{dt} = \frac{\alpha E_T}{1 + \beta E_T} - MN \quad \text{set } E_T = \phi N \quad \text{and} \quad \begin{aligned} \alpha_p &= \alpha \phi \\ \beta_p &= \beta \phi \end{aligned}$$

$$\frac{dN}{dt} = \frac{\alpha_p N}{1 + \beta_p N} - MN$$

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right)$$

Reason for writing the
BH-SRR as above

Here's steepness (Mangel et al 2010, 2013)

$$h = \frac{\frac{\alpha_p}{M}}{4 + \frac{\alpha_p}{M}}$$

Really sweet interpretation

Thing #7: Clear Reference Points Come From the Production Model

Thing #7: Clear Reference Points Come From the Production Model

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right)$$

Thing #7: Clear Reference Points Come From the Production Model

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right)$$

$$\frac{F_{MSY}}{M} = \sqrt{\frac{4h}{1-h}} - 1$$

Thing #7: Clear Reference Points Come From the Production Model

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right)$$

$$\frac{F_{MSY}}{M} = \sqrt{\frac{4h}{1-h}} - 1$$

$$\frac{N_{MNP}}{N_0} = \frac{\sqrt{\frac{4h}{1-h}} - 1}{\frac{4h}{1-h} - 1}$$

Thing #7: Clear Reference Points Come From the Production Model

$$N_0 = \frac{1}{\beta_p} \left(\frac{\alpha_p}{M} - 1 \right)$$

$$\frac{F_{MSY}}{M} = \sqrt{\frac{4h}{1-h}} - 1$$

$$\frac{N_{MNP}}{N_0} = \frac{\sqrt{\frac{4h}{1-h}} - 1}{\frac{4h}{1-h} - 1}$$


$$SPR_{MSY} = \sqrt{\frac{1-h}{4h}}$$

Thing #7: Clear Reference Points Come From the Production Model

An Example

Thing #7: Clear Reference Points Come From the Production Model

An Example



Design considerations for PFMC
groundfish stock assessments and
reference points

Martin Dorn



December 6, 2016
Groundfish Productivity Workshop
Seattle, WA

NOAA
FISHERIES
Alaska Fisheries Science Center

PACIFIC COAST GROUND FISH FISHERY MANAGEMENT PLAN

FOR THE CALIFORNIA, OREGON, AND
WASHINGTON GROUND FISH FISHERY

PACIFIC FISHERY MANAGEMENT COUNCIL
7700 NE AMBASSADOR PLACE, SUITE 101
PORTLAND, OR 97220
(503) 820-2280
(866) 806-7204
WWW.PCOUNCIL.ORG

AUGUST 2016

Thing #7: Clear Reference Points Come From the Production Model

An Example



Design considerations for PFMC
groundfish stock assessments and
reference points

Martin Dorn




December 6, 2016
Groundfish Productivity Workshop
Seattle, WA

PACIFIC COAST GROUND FISH FISHERY MANAGEMENT PLAN

FOR THE CALIFORNIA, OREGON, AND
WASHINGTON GROUND FISH FISHERY

PACIFIC FISHERY MANAGEMENT COUNCIL
7700 NE AMBASSADOR PLACE, SUITE 101
PORTLAND, OR 97220
(503) 820-2280
(866) 806-7204
WWW.PCOUNCIL.ORG

AUGUST 2016



Thing #7: Clear Reference Points Come From the Production Model

An Example



PACIFIC COAST GROUND FISH FISHERY MANAGEMENT PLAN

FOR THE CALIFORNIA, OREGON, AND WASHINGTON GROUND FISH FISHERY

PACIFIC FISHERY MANAGEMENT COUNCIL
7700 NE AMBASSADOR PLACE, SUITE 101
PORTLAND, OR 97220
(503) 820-2280
(866) 806-7204
WWW.PCOUNCIL.ORG

AUGUST 2016



- Ongoing work (EJ Dick, Nick Grunloh, Mangel): W. How much of a difference will it make if a SRR is chosen that hits the reference points?

Thing #7: Clear Reference Points Come From the Production Model

An Example



PACIFIC COAST GROUND FISH FISHERY MANAGEMENT PLAN

FOR THE CALIFORNIA, OREGON, AND WASHINGTON GROUND FISH FISHERY

PACIFIC FISHERY MANAGEMENT COUNCIL
7700 NE AMBASSADOR PLACE, SUITE 101
PORTLAND, OR 97220
(503) 820-2280
(866) 806-7204
WWW.PCOUNCIL.ORG

AUGUST 2016



- Ongoing work (EJ Dick, Nick Grunloh, Mangel): W. How much of a difference will it make if a SRR is chosen that hits the reference points?
- How good an approximation is the production model?

*Thing #8: Setting Steepness Equal to 1 Is About As Non-Conservative
As You Can Get*

*Thing #8: Setting Steepness Equal to 1 Is About As Non-Conservative
As You Can Get*

$$h = \frac{\frac{\alpha_p}{M}}{4 + \frac{\alpha_p}{M}}$$

*Thing #8: Setting Steepness Equal to 1 Is About As Non-Conservative
As You Can Get*

$$h = \frac{\frac{\alpha_p}{M}}{4 + \frac{\alpha_p}{M}}$$

Infinitely productive stock OR
No natural mortality

*Thing #8: Setting Steepness Equal to 1 Is About As Non-Conservative
As You Can Get*

$$h = \frac{\frac{\alpha_p}{M}}{4 + \frac{\alpha_p}{M}} \quad \text{Infinitely productive stock OR} \\ \text{No natural mortality}$$

$$\frac{F_{MSY}}{M} = \sqrt{\frac{4h}{1-h}} - 1 \quad F_{MSY} \rightarrow \infty \text{ as } h \rightarrow 1$$

*Thing #8: Setting Steepness Equal to 1 Is About As Non-Conservative
As You Can Get*

$$h = \frac{\frac{\alpha_p}{M}}{4 + \frac{\alpha_p}{M}} \quad \text{Infinitely productive stock OR} \\ \text{No natural mortality}$$

$$\frac{F_{MSY}}{M} = \sqrt{\frac{4h}{1-h}} - 1 \quad F_{MSY} \rightarrow \infty \text{ as } h \rightarrow 1$$



*Thing #9: The Production Model Predicts Results from Age-Structured
Ones Very Well*

Thing #9: The Production Model Predicts Results from Age-Structured Ones Very Well

Hierarchical Bayesian estimation of recruitment parameters and reference points for Pacific rockfishes (*Sebastes* spp.) under alternative assumptions about the stock-recruit function

Robyn E. Forrest, Murdoch K. McAllister, Martin W. Dorn, Steven J.D. Martell, and Richard D. Stanley


CJFAS 65:286 (2008)

Thing #9: The Production Model Predicts Results from Age-Structured Ones Very Well

Hierarchical Bayesian estimation of recruitment parameters and reference points for Pacific rockfishes (*Sebastes* spp.) under alternative assumptions about the stock–recruit function

Robyn E. Forrest, Murdoch K. McAllister, Martin W. Dorn, Steven J.D. Martell, and Richard D. Stanley

CJFAS 65:286 (2008)





Thing #9: The Production Model Predicts Results from Age-Structured Ones Very Well

Hierarchical Bayesian estimation of recruitment parameters and reference points for Pacific rockfishes (*Sebastes* spp.) under alternative assumptions about the stock-recruit function

Robyn E. Forrest, Murdoch K. McAllister, Martin W. Dorn, Steven J.D. Martell, and Richard D. Stanley

CJFAS 65:286 (2008)




Thing #9: The Production Model Predicts Results from Age-Structured Ones Very Well

Common Name	Scientific Name	Citation
	<i>Scorpaenichthys</i>	
Cabazon	<i>marmoratus</i>	Cope and Key 2009
Canary Rockfish	<i>Sebastes pinniger</i>	Stewart 2009
Cowcod	<i>Sebastes levis</i>	Dick et al. 2009
Darkblotched		Wallace and Hamel
Rockfish	<i>Sebastes crameri</i>	2009
Greenstriped		
Rockfish	<i>Sebastes elongatus</i>	Hicks et al. 2009
Lingcod	<i>Ophiodon elongatus</i>	Hamel et al. 2009
Splitnose		
Rockfish	<i>Sebastes diploproa</i>	Gertseva et al. 2009
Arrowtooth		Kaplan and Helser
Flounder	<i>Atheresthes stomias</i>	2007
Black Rockfish,		
North	<i>Sebastes melanops</i>	Wallace et al. 2007
Black Rockfish,		
South	<i>Sebastes melanops</i>	Sampson 2007
Blue Rockfish	<i>Sebastes mystinus</i>	Key et al. 2008
Chilipepper		
Rockfish	<i>Sebastes goodei</i>	Field 2008
		Gertseva and
Longnose Skate	<i>Raja rhina</i>	Schirripa 2008

Thing #9: The Production Model Predicts Results from Age-Structured Ones Very Well


Common Name	Scientific Name	Citation
	<i>Scorpaenichthys</i>	
Cabazon	<i>marmoratus</i>	Cope and Key 2009
Canary Rockfish	<i>Sebastes pinniger</i>	Stewart 2009
Cowcod	<i>Sebastes levis</i>	Dick et al. 2009
Darkblotched		Wallace and Hamel
Rockfish	<i>Sebastes crameri</i>	2009
Greenstriped		
Rockfish	<i>Sebastes elongatus</i>	Hicks et al. 2009
Lingcod	<i>Ophiodon elongatus</i>	Hamel et al. 2009
Splitnose		
Rockfish	<i>Sebastes diploproa</i>	Gertseva et al. 2009
Arrowtooth		Kaplan and Helser
Flounder	<i>Atheresthes stomias</i>	2007
Black Rockfish,		
North	<i>Sebastes melanops</i>	Wallace et al. 2007
Black Rockfish,		
South	<i>Sebastes melanops</i>	Sampson 2007
Blue Rockfish	<i>Sebastes mystinus</i>	Key et al. 2008
Chilipepper		
Rockfish	<i>Sebastes goodei</i>	Field 2008
		Gertseva and
Longnose Skate	<i>Raja rhina</i>	Schirripa 2008



Mangel et al CJFAS (2013)

Thing #9: The Production Model Predicts Results from Age-Structured Ones Very Well

Common Name	Scientific Name	Citation
	<i>Scorpaenichthys</i>	
Cabazon	<i>marmoratus</i>	Cope and Key 2009
Canary Rockfish	<i>Sebastes pinniger</i>	Stewart 2009
Cowcod	<i>Sebastes levis</i>	Dick et al. 2009
Darkblotched		Wallace and Hamel
Rockfish	<i>Sebastes crameri</i>	2009
Greenstriped		
Rockfish	<i>Sebastes elongatus</i>	Hicks et al. 2009
Lingcod	<i>Ophiodon elongatus</i>	Hamel et al. 2009
Splitnose		
Rockfish	<i>Sebastes diploproa</i>	Gertseva et al. 2009
Arrowtooth		Kaplan and Helser
Flounder	<i>Atheresthes stomias</i>	2007
Black Rockfish,		
North	<i>Sebastes melanops</i>	Wallace et al. 2007
Black Rockfish,		
South	<i>Sebastes melanops</i>	Sampson 2007
Blue Rockfish	<i>Sebastes mystinus</i>	Key et al. 2008
Chilipepper		
Rockfish	<i>Sebastes goodei</i>	Field 2008
		Gertseva and
Longnose Skate	<i>Raja rhina</i>	Schirripa 2008




Mangel et al CJFAS (2013)





*Thing #10: Faster Growth Makes the Age-Structured Model More and More
Like the Production Model*

*Thing #10: Faster Growth Makes the Age-Structured Model More and More
Like the Production Model*



*Thing #10: Faster Growth Makes the Age-Structured Model More and More
Like the Production Model*



*Thing #11, Bonus Track: If You Don't Know What Determines Recruitment,
The Right Prior Is Uniform, Tied Down at 0.2 and 1*

*Thing #11, Bonus Track: If You Don't Know What Determines Recruitment,
The Right Prior Is Uniform, Tied Down at 0.2 and 1*

How to be Wrong

*Thing #11, Bonus Track: If You Don't Know What Determines Recruitment,
The Right Prior Is Uniform, Tied Down at 0.2 and 1*

How to be Wrong

$$h = 1$$

*Thing #11, Bonus Track: If You Don't Know What Determines Recruitment,
The Right Prior Is Uniform, Tied Down at 0.2 and 1*

How to be Wrong

$$h = 1 \quad \text{means} \quad \Pr\{R(0.2B_0) = R_0\} = 1$$

*Thing #11, Bonus Track: If You Don't Know What Determines Recruitment,
The Right Prior Is Uniform, Tied Down at 0.2 and 1*

How to be Wrong

$$h = 1 \quad \text{means} \quad \Pr\{R(0.2B_0) = R_0\} = 1$$

With certainty, recruitment at 20% of unfished biomass is unfished recruitment. You know a lot about what determines recruitment.


*Thing #11, Bonus Track: If You Don't Know What Determines Recruitment,
The Right Prior Is Uniform, Tied Down at 0.2 and 1*

How to be Wrong

$$h = 1 \quad \text{means} \quad \Pr\{R(0.2B_0) = R_0\} = 1$$

With certainty, recruitment at 20% of unfished biomass is unfished recruitment. You know a lot about what determines recruitment.

How to Not be Wrong



Thing #11, Bonus Track: If You Don't Know What Determines Recruitment, The Right Prior Is Uniform, Tied Down at 0.2 and 1


How to be Wrong

$$h = 1 \quad \text{means} \quad \Pr\{R(0.2B_0) = R_0\} = 1$$

With certainty, recruitment at 20% of unfished biomass is unfished recruitment. You know a lot about what determines recruitment.

How to Not be Wrong

If recruitment at 20% of unfished biomass can take any value between 20% of unfished recruitment and unfished recruitment, steepness ranges between 0.2 and 1.0 (tied down at both ends by biology).




Thing #11, Bonus Track: If You Don't Know What Determines Recruitment, The Right Prior Is Uniform, Tied Down at 0.2 and 1

How to be Wrong


$$h = 1 \quad \text{means} \quad \Pr\{R(0.2B_0) = R_0\} = 1$$

With certainty, recruitment at 20% of unfished biomass is unfished recruitment. You know a lot about what determines recruitment..



How to Not be Wrong

If recruitment at 20% of unfished biomass can take any value between 20% of unfished recruitment and unfished recruitment, steepness ranges between 0.2 and 1.0 (tied down at both ends by biology).



Michielsens and McAllister.
2004. CJFAS 61:1032-1047

Some Citations

2010 Mangel, M., Brodziak, J.K.T., and G. DiNardo. Reproductive ecology and scientific inference of steepness: a fundamental metric of population dynamics and strategic fisheries management. *Fish and Fisheries* 11:89-104. **Erratum**

2013 Mangel, M., MacCall, A.D., Brodziak, J., Dick, E.J., Forrest, R.E., Pourzand, R., and S. Ralston. A perspective on steepness, reference points, and stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* 70:930-940

2014 Brodziak, J., Mangel, M. and C-L Sun Stock-recruitment resilience of North Pacific striped marlin based on reproductive ecology. *Fisheries Research*, <http://dx.doi.org/10.1016/j.fishres.2014.08.008>



Options for Moving Forward

Do Not Fix Steepness and Mortality Rate

Options for Moving Forward

Do Not Fix Steepness and Mortality Rate

Replace the BH-SRR by a SRR that Avoids the Problem

Do Not Fix Steepness and Mortality Rate

When will data be informative?

Use Simulation Methods to determine what kinds of data are necessary so that steepness and natural mortality can be estimated in the stock assessment

Do Not Fix Steepness and Mortality Rate

When will data be informative?

Use Simulation Methods to determine what kinds of data are necessary so that steepness and natural mortality can be estimated in the stock assessment

Already started:

Lee, H.-H. et al. 2011. Estimating natural mortality within a fisheries stock assessment model: An evaluation using simulation analysis based on twelve stock assessments. Fisheries Research 109: 89-94.

Lee, H.-H. et al. 2012. Can steepness of the stock–recruitment relationship be estimated in fishery stock assessment models? Fisheries Research 125-126: 254-261.

Replace the BH-SRR by a SRR that Avoids the Problem

An example: Maynard Smith/Shepherd model

$$\frac{dB}{dt} = \frac{\alpha_p B}{1 + \beta B^{\frac{1}{n}}} (M + F) B$$

$$\frac{F_{MSY}}{M} = \frac{\frac{\alpha}{M}(1-n) + \sqrt{\left(\frac{\alpha}{M}\right)^2 (1-n)^2 + 4\frac{\alpha n}{M}}}{2} - 1$$

There is still an undetermined parameter for the analysis -- the data can tell us something! -and if not we need to integrate over the potential range of n